

DB068: Cbl (C14)

Background:

The proto-oncogene c-Cbl was initially identified as the cellular homologue of v-Cbl oncogene that induces pre-B cell lymphomas and myeloid leukemias in mice (1&2). In more recent studies Cbl has been shown to be a negative regulator of tyrosine kinase signaling (2-4). The ubiquitin ligase activity of Cbl leads to the degradation of tyrosine kinases, thus attenuating the signal of receptors (3&4). Targets of Cbl include activated protein tyrosine kinases belonging to the Src and Syk/Zap-70 families (3). An additional mechanism to attenuate receptor signaling is thought to be achieved by Cbl's interaction with downstream targets of tyrosine kinases, such as PI-3K and Vav (1&3).

Origin:

Cbl ($\overline{C}14$) is provided as an affinity purified rabbit polyclonal antibody, raised against a peptide mapping to the carboxy terminal domain of human Cbl p120.

Product Details:

Each vial contains 200 μ g/ml of affinity purified rabbit IgG, Cbl (C14) DB068, in 1 ml PBS containing 0.1 % sodium azide and 0.2% gelatin.

Competition Studies:

A blocking peptide is also available, DB068P, for use in competition studies. Each vial contains 100 µg of peptide in 0.5 ml PBS with 0.1% sodium azide and 100 µg BSA.

Specificity:

Cbl (C14) is recommended to detect mouse, rat, and human Cbl by western blotting, immunoprecipitation, and immunohistochemistry. Recommended western blotting starting dilution 1:200.

Storage:

Store this product at 4° C, do not freeze. The product is stable for one year from the date of shipment.

References:

- 1. Liu YC, Altman A. 1998. Cbl: complex formation and functional implications. Cell Signal 10(6):377-385.
- 2. Miyake S, Lupher ML, Andoniou CE, Lill NL, Ota S, Douillard P, Rao N, Band H. 1997. The Cbl protooncogen product: from an enigmatic oncogene to center stage of signal transduction. Crit Rev Oncog 8(2-3):189-218.
- 3. Rao N, Dodge I, Band H. 2002. The Cbl family of ubiquitin ligases: critical negative regulators of tyrosine kinase signaling in the immune system. J Leukoc Biol 71(5):753-763.
- 4. Sanjay A, Horne WC, Baron R. 2001. The Cbl family: ubiquitin ligases regulating signaling by tyrosine kinase. Sci STKE 2001(110):PE40.