

DB012: cyclin A (C19)

Background:

During each cell cycle cyclins undergo periodic accumulation and destruction. As key regulators of the cell cycle the cyclins control important transitions by activating Cdks (1,2). Cyclin A regulates at least two cdks, Cdc2 p34 and Cdk2, and is believed to be necessary for progression through S phase and is active during the G2-M transition (1&3,4). Cyclin B1 also associates with Cdc2 p34 and this complex is requisite for the G2-M progression. The Cyclin B1/Cdc2 p34 complex is known as MPF, M-phase kinase, maturation-promoting factor, or M-phase specific histone kinase, is widely accepted as the trigger of mitosis in all organisms (5-9).

Origin:

Cyclin A (C19) is provided as an affinity purified rabbit polyclonal antibody, raised against a peptide mapping to the carboxy terminus of mouse cyclin A.

Product Details:

Each vial contains 200 μg/ml of affinity purified rabbit IgG, cyclin A (C19) DB012, in 1 ml PBS containing 0.1 % sodium azide and 0.2% gelatin.

Competition Studies:

A blocking peptide is also available, DB012P, for use in competition studies. Each vial contains 100 µg of peptide in 0.5 ml PBS with 0.1% sodium azide and 100 µg BSA.

Specificity:

Cyclin A (C19) DB012 reacts with cyclin A of mouse, rat, and human origin by western blotting, immunoprecipitation and immunohistochemistry.

Storage:

Store this product at 4° C, do not freeze. The product is stable for one year from the date of shipment.

References:

- Bandara LR, Adamczewski JP, Zamanian M, Poon RY, Hunt T, Thangue NB. 1992. Cyclin A recruits p33cdk2 to the cellular transcription factor DRTF1. J Cell Sci Suppl 16: 77-85.
- 2. Desdouets C, Sobczak-Thepot J, Murphy M, Brechot C. 1995. Cyclin A: Function and expression during cell proliferation. Prog Cell Cycle Res 1:115-123.
- 3. Brechot C. 1993. Oncogenic activation of cyclin A. Curr Opin Genet Dev 3(1): 11-18.
- 4. Goldstone S, Pavey S, Forrest A, Sinnamon J, Gabrielli B. 2001. Cdc25-dependent activation of cyclin A/cdk2 is blocked in G2 phase arrested cells independently of ATM/ATR. Oncogene 20(8): 921-932.
- 5. Liu D, Liao C, Wolgemuth DJ. 2000. A role for cyclin A1 in the activation of MPF and G2-M transition during meiosis of male germ cells in mice. Dev Biol 224(2): 338-400.
- 6. Hoffman I, Clarke PR, Marcotte MJ, Karsenti E, Draetta G. 1993. Phosphorylation and activation of human cdc25-C by cdc2-cyclin B and its involvement in the self-amplification of MPF at mitosis. EMBO J 12(1): 53-63.
- 7. Kishimoto T, Okumura E. 1997. In vivo regulation of the entry into M-phase: initial activation and nuclear translocation of cyclin B/Cdc2. Prog Cell Cycle Res 3:241-249.
- 8. Baratte B, Meijer L, Galaktionov K, Beach D. 1992. Screening for antimitotic compounds using the cdc25 tyrosine phosphatase, an activator of the mitosis-inducing p34cdc2/cyclin Bcdc13 protein kinase. Anticancer Res 12(3): 873-880.
- 9. Thron CD. 1994. Theoretical dynamics of the cyclin B-MPF system: a possible role for p13suc1. Biosystems 32(2):97-109.